

Effect of Dietary Supplementation of Neem oil (Azadirachtaindica) on the Growth performance and Nutrient digestibility of weaned rabbits

¹Oluwafemi R. A, ²Oluwayinka E. O, ³Alagbe J. O

^{1,2,3} Department of Animal Science, Faculty of Agricultural Sciences, University of Abuja, FCT-Abuja. Nigeria,

Article Information

Received : 29 Sep 2020
Revised : 15 Oct 2020
Accepted : 29 Oct 2020
Published : 29 Oct 2020

Abstract— The objective of the present study was to determine the effect of dietary supplementation of neem oil (Azadirachtaindica) on the growth performance and nutrient digestibility of weaned rabbits. A total of 50 weaned male cross bred rabbits between 5-6 weeks with an average weight of 565.4g-566.8g were divided into five dietary groups of ten (10) weaned rabbits each in a completely randomized design. The dietary treatments include a control, T1 (basal) diet with no neem oil (NOL), T2, T3, T4 and T5 were fed basal diet supplemented with NOL at 0.1%, 0.2%, 0.3% and 0.4% respectively. Feed and water were offered ad libitum throughout the experiment which lasted for 12 weeks. The data obtained was used to evaluate the growth performance: weight gain (WG), feed intake (FI), feed conversion ratio (FCR), mortality and nutrient digestibility (dry matter, crude protein, crude fibre and ether extract). WG, ADFI, TFI were significantly ($P < 0.05$) different among the treatments. T5 had the highest weight gain (755.90 g) followed by T4(734.0g), T3(705.90g), T2(705.0g) and T1(621.80g) respectively. Highest mortality was recorded in T1 (2%) followed by T2 (1%), none was recorded in T3, T4 and T5. Neem oil significantly influenced ($P < 0.05$) all the parameters measured. It could be concluded that neem oil contains some essential nutrients and bioactive chemicals and could be supplemented in the diet of rabbit at 0.4 % without any deleterious effect on the general performance of the animal.

Corresponding Author:

Oluwayinka E. O Email:
olaoluwayinkamail@gmail.com

Keywords: Azadirachtaindica, neem seeds, nutrients, performance, Rabbits.

Copyright © 2020: Oluwafemi R. A, Oluwayinka E. O and Alagbe J. O. This is an open access distribution, and reproduction in any medium, provided Access article distributed under the Creative Commons Attribution License the original work is properly cited License, which permits unrestricted use.

Citation: Oluwafemi R. A, Oluwayinka E. O and Alagbe J. O. “Effect of Dietary Supplementation of Neem oil (Azadirachtaindica) on the Growth performance and Nutrient digestibility of weaned rabbits”, Journal of Science, Computing and Engineering Research, 1(4), 100-105, September-October 2020.

I. INTRODUCTION

With so many research carried out to find endogenous feed alternatives for rabbit production as the high cost of raw materials that are used for commercial feed increase the production cost., there is also an urgent need to embark on scientific findings on livestock's natural growth promoter which will ensure increasing growth rate, enormous availability of meat and other high value by-products for the entire populace at relatively conservative cost. Producers use growth promoters to increase growth rates and improve overall efficiency and product quality without leaving any toxic residue in the body system of livestock. Their inclusion in feedstuffs should be designed so as to feature a pharmacological characteristic that enhance the immunity of the animal and to help in minimizing the use of the conventional antibiotics in prevention and treatment of diseases of livestock. The conventional artificial growth promoters are known to have deleterious effect on human

who are the secondary consumer of residues of artificial growth promoter in the body of livestock (Sinniah, 1981). Neem belongs to the kingdom: Plantae; Division:

Magnoliophyta; Order: Sapindales; Family: Meliaceae; Genus: Azadirachta; Species: indica., It is a tropical evergreen related to mahogany. Native to east India and Burma, it grows in much of Southeast Asia and West Africa; a few trees have recently been planted in the Caribbean and several Central American countries, including México. The name Azadirachtaindica is derived from a Persian term “Axaddarakth” (free tree). In Ayurveda it is known as the ‘Arishta’, which means “relieving sickness” in Sanskrit. It is a medium sized or large evergreen tree with irregular rounded crown, attaining a height of 14m-20m. It is a hardy tree that grows well in sandy, stony shallow soil, and is tolerant to alkaline, saline and acidic soil and it grows well on black cotton soil (Patnaik, 1993).

Neem is ubiquitous in Northern Nigeria. The Neem tree popularly referred to in Hausa language as Dogonyaro is a tree in the mahogany family with broad dark brown stem and widely spread branches. According to Subbalakshmi et al., (2012), all parts of neem like seeds, flowers, bark and leaves are beneficial due to their medicinal properties. Research has shown that neem will boost the immune system by stimulating the production of T-cells when challenged with

Effect of Dietary Supplementation of Neem oil (*Azadirachta indica*) on the Growth performance and Nutrient digestibility of weaned rabbits

Available at <https://jscer.org>

infections (Upadhyay, 1990). The role of medicinal plants in disease prevention or control has been attributed to antioxidant properties of their constituents, usually associated to a wide range of amphipathic molecules, broadly termed polyphenolic compounds (Demirayet et al., 2009). The bark of the neem has been reported to have higher phenolic and antioxidant activity compared to the leaf (Ghimeray et al., 2009; Olabinri et al., 2009).

Neem oil, bark and leaf extracts have been therapeutically used as folk medicine to control diseases like leprosy, intestinal helminthiasis, respiratory disorders, constipation and skin infections (Biswas et al., 2002). The neem tree contains more than 100 bioactive ingredients and the most important bioactive compound is azadirachtin (Nahak and Sahu, 2010). The Neem leaves, neem oil and de-oiled neem seed cake are used as animal feeds (Ogbuwuet al., 2010a). The neem leaves contain appreciable amounts of proteins, minerals, carotene and adequate amount of trace minerals (Ogbuwuet al., 2010). Neem tree as one of the most researched tree in the world has attracted world-wide prominence due to its vast range of medicinal properties like antibacterial, antiviral, antifungal, antiprotozoal, hepatoprotective and other various properties without showing any adverse effect (Kale et al., 2003).

The compounds in neem have been divided into two major classes; isoprenoids and others (Singh et al., 1996). The isoprenoids include diterpenoids and triterpenoids containing protomeliacins, limonoids, azadirone and its derivatives, gedunin and its derivatives, vilasinin type of compounds and Csecomeliacins such as nimbin, salanin and azadirachtin. The none-isoprenoids include proteins (amino acids) and carbohydrates (polysaccharides), sulphurous compounds, polyphenolics such as flavonoids and their glycosides, dihydrochalcone, coumarin and tannins, aliphatic compounds and several fatty acids (dodecanoic, tetradecanoic, elcosanic, etc). (Zengin et al., 2016a, Zengin et al., 2016b).

II. MATERIAL AND METHODS

A. Experimental Site

The study was carried out at University of Abuja Teaching and Research farm, Airport road, Abuja., in Gwagwalada area council of the Federal Capital Territory, Abuja. Gwagwalada is situated in the North central zone of Nigeria., Lying at the latitude N 9.0765 and longitude E 7.3986 at an average elevation of 476m above sea level.

B. Collection And Processing Of Neem Oil (No)

Neem seeds were collected from Gwagwalada, Abuja and identified at the department of biological sciences, University of Abuja, Nigeria. The seeds of Neem were separated from the seed coats manually and sundried for 2 weeks. The dried seeds were granulated into coarse particles using a blender (Model Ap-DKL, Samsung). Oil was extracted using the soxhlet extraction method; it was later poured into a well labeled container for further analysis.

C. Experimental Animals And Their Management

Fifty (50) apparently healthy, cross bred weaned male rabbits with average initial body weight of 565.4g-566.8g were used for the study and were randomly allotted into Five Treatments with ten (10) rabbits per treatment designated as treatment 1, 2, 3, 4 and 5 in a Completely Randomized Design (CRD), animals were kept in an all wired hutch measuring 35 × 35 × 55cm (width × length × height).

All treatments have 5 replicates with two (2) rabbits per replicate. After 14 days of acclimatization, all rabbits were fed diets corresponding to their treatments and given prophylactic treatment with broad-spectrum medication (Kepromec®) against endoparasites and helminthes infestation before the commencement of the experiment. Feed and water was given ad libitum and all other management practices were strictly adhered to.

D. Experimental Diets

Basal were formulated to meet the nutritional requirement for rabbits according to NRC (1977).

- Treatment 1 – Basal diet + 0 % NOL
- Treatment 2 – Basal diet + 0.1 % NOL
- Treatment 3 – Basal diet + 0.2 % NOL
- Treatment 4 – Basal diet + 0.3 % NOL
- Treatment 5 – Basal diet + 0.4 % NOL

E. Measurements

Daily feed consumption (g) = Amount of feed Supplied – Amount of feed left

Feed conversion ratio (FCR) = $\frac{\text{Dry Matter Intake (g)} \times 100}{\text{Live weight gain}}$

Average daily gain (ADG) = $\frac{\text{Final body weight} - \text{Initial body weight}}{\text{Total days of the experiment}}$

F. Digestibility Trial

At the last week of the experiments, fifteen (15) rabbits (3 per treatment) were randomly selected and were housed individually in metabolic cages which allow separation of faeces and urine. Each cage was provided with manual feeder and drinker. The digestibility study lasted seven days, average feed consumption 48.60 – 49.04g was accurately determined during the whole experiment period and faeces were collected for seven days as a collection period (Perez et al., 1995).

The collection was performed at approximately 09:00h each morning before the next daily ration was provided. The rabbits are fed measured amounts of feed and the left over feed and faeces voided are carefully measured .5% of faeces voided were dried at 800C in an oven for 24 hours. The dried faecal samples were stored in air-tight container for chemical analysis. This can be represented using the equation;

Digestibility(%) = $\frac{C_{\text{feed}} - C_{\text{faeces}}}{C_{\text{feed}}} \times 100$

Where C_{feed} and C_{faeces} refer to the amount of feed eaten and faecal excreted respectively. The proximate composition

Effect of Dietary Supplementation of Neem oil (*Azadirachtaindica*) on the Growth performance and Nutrient digestibility of weaned rabbits

Available at <https://jscer.org>

of the diets and faecal samples were determined according to AOAC (1990).

G. Statistical Analysis

All data were subjected to one -way analysis of variance (ANOVA) using SPSS (18.0) and significant means were separated using Duncan multiple range tests (Duncan, 1955). Significant was declared if $P \leq 0.05$.

III. RESULTS AND DISCUSSION

A. Proximate composition of experimental diet

Table I shows the proximate composition of experimental diet. The proximate components contained crude protein (18.22 %), crude fibre (13.22 %), ether extract (3.20 %), ash (6.15 %) and energy (2566.5 kcal/kg). The crude protein, crude fibre and energy values reported in this experiment is in agreement with the findings of Ahmed et al. (2018); Alagbe and Oluwafemi (2019) but contrary to the reports of Ahmed et al. (2019) when thyme oil was fed to growing rabbits.

However, all values were within the nutritional requirement of growing rabbits according to NRC (1977). Adequate intake of dietary fibre lowers the serum cholesterol level, risk of coronary heart disease, constipation and colon and breast cancer (Fashola, 2011; Alagbe, 2019; Olanipekun et al., 2016). Ash content gives an indication of the amount of minerals present in a feed, which are important in many biochemical reactions functioning as co-enzyme and aid physiological functioning of the major metabolic processes in the body (Ojewuyi et al., 2014).

Table I. Chemical composition of experimental diet

Materials	Quantity (Kg)
Maize	30.0
Wheat offal	20.0
Soya meal	16.25
Groundnut cake	10.0
Palm kernel meal	20.0
Bone meal	2.00
Limestone	1.00
Lysine	0.01
Methionine	0.01
*Premix	0.25
Salt	0.25
Total	100.0
Calculated analysis	
Crude protein (%)	17.22
Crude fibre (%)	13.20

Ether extract (%)	3.02
Ash (%)	6.15
Energy (Kcal/kg)	2566.5

B. Effect of different levels of neem (*Azadirachtaindica*) oil on the performance of weaned rabbits

Table II revealed the performance characteristics of weaned rabbits graded level of neem oil. Initial body weight (IBW), final body weight (FBW), weight gain (WG), average daily weight gain (ADWG), average weekly weight gain (AWWG) and total feed intake (TFI) ranged between 566.20 – 566.80 g, 1188.40 – 1322.10 g, 6.21 – 755.9 g, 8.88 – 10.80 g, 62.18 – 75.60 g and 2900.1 – 3433.1 g respectively. All the values were higher in T4 and T5, intermediate in T2 and T3 and lowest in T1 ($P < 0.05$). FCR value ranged between 4.50 – 5.50 and it was significantly different among the treatments ($P < 0.05$). Mortality were recorded in T1 and T2 ($P < 0.05$), none were recorded in the other treatments.

The higher weight gains in T4 and T5 could be attributed to the presence of bioactive chemicals or phytochemicals in neem oil. According to Oluwafemi *et al.* (2020) phytochemicals performs multiple biological activities such as anti-inflammatory (Hyun *et al.*, 2018; Alagbe *et al.*, 2020), antibacterial (Tomoriet *et al.*, 2007), antioxidant (Manita and Gaurav, 2020), immunostimulatory (Kavita *et al.*, 2014), antifungal, hepatoprotective and cardiovascular effects (Joy *et al.*, 2019; Olafadehan *et al.*, 2020), antiviral (Olafadehan *et al.*, 2020). Bioactive chemicals include: tannins, saponins, flavonoids, alkaloids, phenols, terpenoids etc. The supplementation of neem oil in the diet of the animals promotes eubiosis, thus enhancing the activities of beneficial bacteria (lactobacilli) and lowering pathogenic bacteria (Alagbe *et al.*, 2020).

This result is in agreement with the findings of Ikyume *et al.* (2019); Cardinal *et al.* (2007); Castellini *et al.* (2007); Abd El-Hady *et al.* (2013) but contrary to the reports of Eiben *et al.* (2008); Musa *et al.* (2020) on feed additives as they affect the fattening performance of rabbits. Mortality was not recorded in T3, T4 and T5; this could be possibly attributed to the presence of phenols, saponins and tannins. According to Hollman (2001), phenols are strong antioxidants capable of scavenging free radicals. Tannins possess antibacterial and antiviral activities (Adisa *et al.*, 2010; Enzo, 2007) while saponins have been suggested to be involved in antimicrobial and anti-inflammatory activities (Cheeke 2000; Alagbe *et al.*, 2019; Akintayo and Alagbe, 2020).

C. Nutrient digestibility of rabbits fed graded levels of neem oil

Nutrient digestibility of rabbits fed graded levels of neem oil is presented in Table 3. Dry matter, crude protein, crude fibre and ether extract digestibility ranged between 75.11 – 89.30 %, 60.22 – 78.90 %, 50.92 – 69.31 % and 67.89 – 70.47 % respectively. There was a significant difference in

Effect of Dietary Supplementation of Neem oil (*Azadirachtaindica*) on the Growth performance and Nutrient digestibility of weaned rabbits

Available at <https://jscer.org>

Table II. Effect of different levels of neem (*Azadirachtaindica*) oil on the performance of weaned rabbits

Parameters	T1	T2	T3	T4	T5	SEM
IBW (g)	566.60	565.40	566.80	566.50	566.20	0.31
FWG (g)	1188.40 ^c	1270.40 ^b	1272.70 ^b	1300.50 ^a	1322.10 ^a	9.98
WG (g)	621.80 ^c	705.00 ^a	705.90 ^b	734.00 ^a	755.90 ^a	2.11
ADWG (g)	8.88 ^d	10.07 ^c	10.08 ^c	10.48 ^b	10.80 ^a	0.87
AWWG (g)	62.18 ^d	70.50 ^c	70.59 ^c	73.40 ^b	75.60 ^a	0.04
TFI (g)	3433.10 ^a	3420.10 ^a	3411.80 ^a	3409.20 ^a	2900.10 ^b	11.76
FCR	5.50 ^a	4.85 ^b	4.83 ^b	4.64 ^c	4.50 ^d	0.02
MORT.	2.00	1.00	-	-	-	0.01

*Means in the same row with different superscripts differ significantly ($P<0.05$) Initial body weight (IBW); final body weight (FBW); weight gain (WG); average daily weight gain (ADWG); average weekly weight gain (AWWG) and total feed intake (TFI)

the parameters measured ($P<0.05$). This result is in agreement with the findings of Omokore and Alagbe (2019); Bassionyet *et al.* (2015); Bölükbaşı *et al.* (2006); Çabuket *et al.* (2003); Gaafaret *et al.* (2014); Shittu *et al.* (2020) when pumpkin (*Cucurbitamoschata*) and black seed (*Nigella sativa*) oils were supplemented in the diets of rabbits but

contrary to the reports of Taiwo *et al.* (2005); Oso *et al.* (2006) when weaned rabbits were fed rice millings waste based diets. Bioactive chemicals in neem oil ensure proper absorption of nutrient among animals in T4 and T5, thus ensuring better feed conversion among animals.

Table III. Nutrient digestibility of rabbits fed graded levels of neem oil

Parameters (%)	T1	T2	T3	T4	T5	SEM
Dry matter	75.11 ^c	79.21 ^b	80.93 ^b	85.33 ^a	89.30 ^a	2.71
Crude protein	60.22 ^c	65.97 ^b	69.03 ^b	71.41 ^a	78.90 ^a	1.98
Crude fibre	50.92 ^b	57.90 ^b	61.88 ^a	68.81 ^a	69.31 ^a	1.02
Ether extract	67.89 ^b	61.80 ^b	67.98 ^a	69.41 ^a	70.47 ^a	2.33

Means in the same row with different superscripts differ significantly ($P<0.05$)

IV. CONCLUSION

Bioactive chemicals in neem oil which may have acted singly or in synergy with one another ensure proper absorption of nutrients among animals in T4 and T5, thus ensuring better feed conversion among animals. *A. indicain* diets of growing rabbits had a positive effect on growth performance, feed conversion ratio and on mortality rate.

REFERENCES

- [1]. Abd El-Hady, A.M., O. A.H. El-Ghalid and A.M. EL-Raffa. (2013). Influence of a herbal feed additives (digestarom®) on productive performance and blood constituents of growing rabbits. *Egyptian J. Anim. Prod.* (2013) 50(1):27-37.
- [2]. A.O.A.C. (2000). Association of Official Analytical Chemists. Official Methods of Analysis 19th Edition Washington, D.C Pages 69-77.
- [3]. Akintayo Balogun Omolere. M and Alagbe, J.O (2020). Probiotics and medicinal plants in poultry nutrition: A review. *United International Journal for Research and Technology*, 2(1): 7-13.
- [4]. Alagbe, J.O., Agubosi, O.C.P., Ajagbe, A.D, Shittu, M.D and Akintayo Balogun, O.M (2020). Performance, haematology

and serum biochemical parameters of growing grass cutters fed *Phyllanthus amarus* and *Piliostigma thonningii* leaf meal mixture as partial replacement for Soya bean meal. *United International Journal for Research and Technology*, 2(1): 14-23.

- [5]. Ahmed A.A. Abdel-Waret ,Eman M.M. Taha , Karl-Heinz Südekun, Jayant Lohakare (2018). Thyme oil inclusion levels in a rabbit ration: Evaluation of productive performance, carcass criteria and meat quality under hot environmental conditions. *Animal Nutrition* 4 (2018) 410e416
- [6] Ahmed E. Abd El-Azeem, Adham A. Al-Sagheer, A.H. Daader and S.M. Bassiony. (2019). Effect of dietary supplementation with betaine, thyme oil and their mixtures on productive performance of growing rabbits. *Zagazig J. Agric. Res.*, 46(3): 816-827.
- [7] Alagbe, J.O and Oluwafemi, R.A. (2019). Growth performance of weaner rabbits fed Noni (*Morinda citrifolia*) and Moringa *olifera* leaf mixture as partial replacement of soya bean meal. *International Journal of Advanced Biological and Biomedical Research*. 7(2): 185-195.
- [8] Alagbe, J.O (2019). Proximate, mineral and phytochemical analysis of *Piliostigma thonningii* stems bark and roots.

Effect of Dietary Supplementation of Neem oil (*Azadirachta indica*) on the Growth performance and Nutrient digestibility of weaned rabbits

Available at <https://jscer.org>

International Journal of Biological, Physical and Chemical Studies, 1(1): 1-7.

- [9] Alagbe, J.O. (2020). Caecal Microbial Population of Growing Grass Cutters (*Thyronoymys Swinderianus*) Fed *Phyllanthus Amarus* and *Pilostigma Thonngii* Leaf Meal Mixture as Partial Replacement for Soya Bean Meal. *Concept of Dairy and Veterinary Sciences*. 3(5): 350 – 355.
- [10] Alagbe, J.O., Shittu, M.D and Eunice Abidemi Ojo (2020). Prospect of leaf extracts on the performance and blood profile of monogastric – A review. *International Journal of Integrated Education*. 3(7): 122-127.
- [11] Bassiony, S., M.M. Elhindawy, AE. Attia and I.E. Ismail (2015). Effect of some bioactive components of essential oils on growing rabbits' performance. *Zagazig J. Agric. Res.*, 42 (5):1171-1182.
- [12] Biswas K., Chattopadhyay I., Banerjee R. K., Bandyopadhyay U. (2002). Biological activities and medicinal properties of Neem (*Azadirachta indica*) *Current Science*.82(11):1336–1345.
- [13] Bölükbaşı, S.C., M.K. Erhan and A. Özkan (2006). Effect of dietary thyme oil and vitamin E on growth, lipid oxidation, meat fatty acid composition and serum lipoproteins of broilers. *S. Afr. J. Anim. Sci.*, 36: 189–196.
- [14] Çabuk, M., A. Alçiçek, M. Bozkurt and N. Imre (2003). Antimicrobial properties of the essential oils isolated from aromatic plants and using possibility as alternative feed additives. *II. Nat. Anim. Nut. Con.*, 18 (20): 184 -187.
- [15] Castellini C., Cardinali R., Rebollar P.G., Dal Bosco A., Jimeno V., Cossu M.E. 2007. Feeding fresh chicory (*Chicoriaintybus*) to young rabbits: Performance, development of gastro-intestinal tract and immune functions of appendix and Peyer's patch. *Anim. Feed Sci. Technol.*, 134, 56-65.
- [16] Cardinali R., Dal Bosco A., Mourvaki E., Moscati L., Scicutella N., Battistacci L., Castellini C. 2007a. Effect of dietary microencapsulated organic and inorganic acids and essential oils on serum innate and caecal fermentation. In: *Proc. Giornate di Coniglicoltura ASIC 2007*, September, Forlì, Italy, 133.
- [17] Demiray S., Pintado M. E. and Castro P.M.L., (2009) Evaluation of phenolic profiles and antioxidant activities of Turkish medicinal plants: *Tiliaargentea*, *Crataegi folium* leaves and *Polygonumbistoa* roots. *World Acad. Science England. Technol.*, 54:312-317.
- [18] Eiben, C., Gippert, T., Gódor-Surmann K and Kustos, K. (2008). Feed additives as they affect the fattening performance of rabbits. *Nutrition and Digestive Physiology*. 9th World Rabbit Congress – June 10-13, 2008 – Verona – Italy
- [19] Gaafar HMA, Ragab AA, El-Reidy KFA (2014) Effect of diet supplemented with pumpkin (*Cucurbitamoschata*) and black seed (*Nigella sativa*) oils on performance of rabbits. In: *Growth performance, blood haematology and carcass traits of growing rabbits. Reports and Opinions.*, volume 16: 52-59.
- [20] Ghimeray A. K., Jin C. W., Ghimire B. K., Cho D. H. (2009). Antioxidant activity and quantitative estimation of azadirachtin and nimbin in *Azadirachta indica* A. Juss grown in foothills of Nepal African Journal of Biotechnology. 8(13):3084–3091.
- [21] Ikyume, T.T., Ogu I.E, Okwori IA, Shaahu DT. (2019). Growth Performance and Apparent Nutrient Digestibility of Grower Rabbits Fed Combinations of Concentrate with Grass and/or Legume Forage. *Journal of Multidisciplinary Research and Reviews*. 1(1): 41-45.
- [22] Gaafar HMA, Ragab AA, El-Reidy KFA (2014) Effect of diet supplemented with pumpkin (*Cucurbitamoschata*) and black seed (*Nigella sativa*) oils on performance of rabbits. In: *Growth performance, blood haematology and carcass traits of growing rabbits. Reports and Opinions*, volume 16: 52-59.
- [23] Shittu, M.D., Adejumo, D.O., Ewuola, E.O., Alaba, O., Alagbe, J.O and Ojebiyi, O.O. (2020). Gut morphometric characteristic and ecological response of broiler starter fed varied levels of protein. *Asian Journal of Animal Science*, 14(1):33-39
- [24] Ghimeray A. K., Jin C. W., Ghimire B. K., Cho D. H. (2009). Antioxidant activity and quantitative estimation of azadirachtin and nimbin in *Azadirachta indica* A. Juss grown in foothills of Nepal African Journal of Biotechnology. 8(13):3084–3091.
- [25] Ikyume, T.T., Ogu I.E, Okwori IA, Shaahu DT. (2019). Growth Performance and Apparent Nutrient Digestibility of Grower Rabbits Fed Combinations of Concentrate with Grass and/or Legume Forage. *Journal of Multidisciplinary Research and Reviews*. 1(1): 41-45.
- [26] Kale, B.P., M.A. Kothekar, H.P. Tayade, J.B. Jaju and M. Mateenuddin. (2003). Effect of aqueous extract of *Azadirachta indica* leaves on hepatotoxicity induced by antitubercular drugs in rats. *Indian J. Pharmacol.*, 35: 177-180.
- [27] Musa, Bashir., Alagbe, J.O., Adegbite Motunrade Betty, Omokore, E.A. (2020). Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2(2):13-21.
- [28] Nahak G., Sahu R. K. (2011). Evaluation of antioxidant activity of flower and seed oil of *Azadirachta indica* A. juss. *Journal of Applied and Natural Science*. 3(1):78–81.
- [29] Ogbuewu, I.P., Odoemelam, V.U., Obikaonu, H.O., Opara, M.N., Emenalom, O.O., Uchegbu, M.C., Okoli, I.C., Esonu, B.O. and Iloeje, M.U. (2010a). The growing importance of neem (*Azadirachta indica* A. Juss) in agriculture, industry, medicine and environment: A Review. *Research Journal of Medicinal Plants*, 3(2): 1-27.
- [30] Ogbuewu, I.P., Okoli, I.C. and Iloeje, M.U. (2010b). Assessment of blood chemistry, weight gain and linear body measurements of pre-puberal buck rabbits fed different levels of Neem (*Azadirachta indica* A. Juss.) leaf

Effect of Dietary Supplementation of Neem oil (*Azadirachta indica*) on the Growth performance and Nutrient digestibility of weaned rabbits

Available at <https://jscer.org>

- meals. *Chilean Journal of Agricultural Research*, 70(3): 515-520.
- [31] Olabinri, B.M., J.A. Adebisi, O.F. Odesomi, P.F. Olabinri and G.E. Adeleke (2009). Experimental classification of the antioxidant capacity of the leaf, stem and root barks of *Magnifera indica* and *Azadirachta indica*. *Afr. J. Biotechnol.*, 8(13): 2968-2972.
- [32] Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O. (2020). Performance, haemato-biochemical parameters of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Advances in Research and Reviews*, 2020, 1:4.
- [33] Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O (2020). Carcass quality, nutrient retention and caeca microbial population of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Journal of Drug Discovery*. 14(33):146-154.
- [34] Oluwafemi, R.A., Isiaka Olawale and Alagbe, J.O. (2020). Recent trends in the utilization of medicinal plants as growth promoters in poultry nutrition- A review. *Research in: Agricultural and Veterinary Sciences*. 4(1): 5-11
- [35] Oso A.O, Bamgbose AM, Isah OA, Olatunji JEN, Mabadeje AT, Alade AA, Oni AO (2006) Performance of weaner rabbits fed rice millings waste based diets. *Journal of Animal and Veterinary Advances* 5: 836-838.
- [36] Omokore, E.O and Alagbe, J.O. (2019). Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits. *International Journal of Academic Research and Development*. 4(3): 97-104.
- [37] Ojewuyi, O.B, Ajiboye, T. O, Adebajo, E. O, Balogun, A, Mohammed, A.O (2014). Proximate composition, phytochemical and mineral contents of young and mature *Polyalthia longifolia* Sonn. leaves Fountain *Journal of Natural and Applied Sciences*: 2014; 3(1): 10 – 19.
- [38] Patnaik, N. (1993). *Garden of life: An introduction to the healing plants of India*. Doubleday, New York pp. 40 – 42.
- [39] Duncan, D.B. (1955). Multiple range and multiple F-test. *Biometrics* 11(1):1-42. National Research Council (1977). *Nutrient requirement of rabbits 9th Rev Edn*, Washington D.C. National Academy Press.
- [40] Subbalkshmi L., Muthukrishnan P., Jeyaraman S. (2012). *Neem products and their Agricultural application*.
- [41] Taiwo A.A, Adejuyigbe AD, Adebawale EA, Oshotan JS, David OO (2005) Performance and nutrient digestibility of weaned rabbits fed forages supplemented with concentrates. *Nigerian Journal of Animal Production*. 32:74-78
- [42] Upadhyay, C. (1990). The medicinal properties of Neem (*Azadirachta indica*) tree. In: *Animal Pharmacology*. Second Edition Longman England.
- [43] Zengin, G., Ceylan, R. O., Guler, G., Carradori, S., Uysal, S., Aktumsek, A. (2016a). Enzyme inhibitory effect and antioxidant properties of *Astragalus lagurus* extract. *Current enzyme inhibition* 12, 177-182.
- [44] Zengin, G., Locatell, M., Carradori, S., Mocan, A. M., Aktumek, A., (2016b). Total phenolics, flavonoids, condensed tannins content of eight *Centaurea* species and their broad inhibitory activities against cholinesterase, tyrosinase, α -amylase and α -glucosidase. *Notulae Botanica Horti Agrobotanici Cluj-Napoca* 44, 195-200